

CLAIMS

1. A method for evaluating renal functions, which comprises measuring the megsin protein in a biological specimen.

2. The method for evaluating renal functions of claim 1, which comprises measuring the megsin protein in a biological specimen, and comparing it with the megsin protein amount in a normal specimen.

3. The method for evaluating renal functions of claim 1, wherein the biological specimen is urine.

4. The method for evaluating renal functions of claim 1, wherein the megsin protein in the biological specimen is measured by an antigen-antibody reaction using an anti-megsin protein antibody.

5. The method for evaluating renal functions of claim 4, wherein the anti-megsin protein antibody is a monoclonal antibody.

6. A reagent for diagnosing renal functions, which comprises the anti-megsin protein antibody.

7. The reagent for diagnosing renal functions of claim 6, wherein the anti-megsin protein antibody is a monoclonal antibody.

8. A granule for detecting megsin protein in a biological specimen, wherein the granule comprises a solid granule to the surface of which an anti-megsin protein antibody is bound.

9. The granule for detecting megsin protein of claim 8, wherein the solid granule is magnetic.

10. The granule for detecting megsin protein of claim 8, wherein the relative density of the solid granule is not smaller than 1.

11. The granule for detecting megsin protein of claim 8, wherein the anti-megsin protein antibody is a monoclonal antibody.

12. A method for detecting megsin protein in a biological specimen, comprising the following steps of:

i) contacting the granule for detecting megsin protein with the biological specimen, said granule comprising a solid granule to the surface of which an anti-megsin protein antibody is bound.

ii) contacting the second anti-megsin protein antibody bound to a marker molecule with said granule for detecting megsin protein to which the biological specimen was contacted. and,

iii) detecting the marker molecule bound to the megsin protein through

Sub
D1

09936883 122101
TOTAT 8889660

D1

SECRET

14. The method for detection of claim 13, wherein the first anti-megsin protein antibody and the second anti-megsin protein antibody are antibodies having different recognition sites.

16. The method for detection of claim 12, wherein the biological specimen is blood.

5 a) magnetic solid granules to which anti-megsin protein antibodies
can be bound,

18. The kit for detecting megsin proteins of claim 17, further
20 comprising an anti-megsin protein antibody to which a marker molecule
is bound.